

On the possibility of testing the Dvali-Gabadadze-Porrati brane-world scenario with orbital motions in the Solar System

Lorenzo Iorio
*Viale Unità di Italia 68, 70125
Bari, Italy
e-mail: lorenzo.iorio@libero.it*

Abstract

Recently it has been suggested to use the perigee of the proposed LARES/WEBER-SAT satellite in order to measure the secular precession which would be induced on such Keplerian orbital element by a weak-field modification of gravity occurring in some Brane-World scenarios put forth by Dvali, Gabadadze and Porrati. This precession, derived for the first time by Lue and Starkman, amounts to $\sim \mp 4 \times 10^{-3}$ milliarcseconds per year. In this paper we show that, according to the recently released EIGEN-CG01C Earth gravity model which combines data from the dedicated CHAMP and GRACE missions, the quite larger systematic errors due to the Newtonian part of the terrestrial gravitational potential would vanish any attempts to detect a so small effect. Improvements in our knowledge of the Earth's gravitational field of, e.g., up to four orders of magnitude in the low degree even zonal harmonics would be required. Also the obtainable observational accuracy in reconstructing the LARES orbit rules out the possibility of measuring the LS effect with the perigee of LARES. The situation is much more favorable in the Solar System scenario. Indeed, the non-Newtonian perihelion advance of Mars, which is the currently best tracked planet, has recently been measured with an accuracy of 1×10^{-2} milliarcseconds over a 92-years time span; the LS precession is $\sim \mp 4 \times 10^{-1}$ milliarcseconds per century. A suitable combination of the perihelia and the nodes of Mars and Mercury, which disentangles the LS effect from the competing larger Newtonian and general relativistic precessions, might allow to reach a 1-sigma 41% level of accuracy.

Keywords: LARES, even zonal harmonics, planets of the Solar System, orbital motion, Brane-World scenario

Contents

1	Introduction	3
2	The Earth-LARES system	4
2.1	The observational sensitivity	4
2.2	The systematic errors of gravitational origin	4
2.2.1	The impact of the even zonal harmonics of the geopotential	5
2.2.2	The linear combination approach	5
2.2.3	The impact of the odd zonal harmonics	8
3	The Earth-Moon system	8
4	The Solar System scenario	9
4.1	The observational sensitivity	9
4.2	The systematic errors	10
5	Conclusions	13
6	Appendix: The impact of the asteroids on the Mars orbital motion	14

1 Introduction

The recently observed acceleration of the Universe could be explained either by invoking a small vacuum energy (or some effective vacuum energy) or by assuming that the Einstein gravity undergoes long-range modifications imitating a small cosmological constant. The second approach seems to be more promising because it is perturbatively stable under quantum corrections. In it the conventional gravity breaks down at very large distances, beyond a certain threshold r_c which is perturbatively stable and is fixed by the observations to a value $\sim 10^{26}$ m. In such a framework there is a class of Brane-World theories which, for $r \ll r_c$, lead to Einstein-Newton gravity up to small corrections [1]. As a consequence of such tiny deviations, a secular advance of the argument of pericentre ω of a test particle freely orbiting a central mass M occurs.

According to the Dvali-Gabadadze-Porrati (DVP) model [2], which is a Brane-World theory with an infinite-volume flat extra spatial dimension, the analytical expression of the pericentre advance over one orbit of one object moving at distance r from a central body of mass M is [3]

$$\Delta\omega = \mp \frac{3\pi\sqrt{2}}{4} \frac{r^{3/2}}{r_c\sqrt{r_g}} \text{ rad orbit}^{-1}, \quad (1)$$

where $r_c = 6 \text{ Gpc}^1 = 1.8514068 \times 10^{23} \text{ km}$ and $r_g = 2GM/c^2$. The Lue-Starkman (LS) secular rate for any object orbiting a central mass along a circular orbit amounts to $\sim \mp 4 \times 10^{-3}$ milliarcseconds per year (mas yr $^{-1}$), or $\mp 4 \times 10^{-4}$ arcseconds per century (″ cy $^{-1}$), only².

The possibility of testing such an effect in the Solar System scenario has preliminarily been investigated for the first time in [3]. In a recent paper [5] it has been suggested, among other things, to use the perigee of the proposed Earth artificial satellite LARES in order to detect the LS effect.

In the present work we analyze in more details such possibilities yielding a realistic evaluation of the error budgets by also including many relevant sources of systematic errors.

¹1 parsec (pc)= 3.085678×10^{13} km. The value $r_c = 6 \text{ Gpc}$ is quoted in [1, 4]

²Note that in [3] $r_c \sim 5 \text{ Gpc}$ is assumed from fitting of both the Cosmic Microwave Background (CMB) and the type IA supernovae (Starkman, G. 2005; personal communication): this yields $5 \times 10^{-4} \text{ ″ cy}^{-1}$ for the magnitude of the LS effect. Its sign is affected by the cosmological branch [3]: it is negative for the standard cosmological branch, and positive for the self-accelerated one.

Table 1: Orbital parameters of LAGEOS, LAGEOS II and LARES.

Orbital element	LAGEOS	LAGEOS II	LARES
semimajor axis a (km)	1.2270×10^4	1.2163×10^4	1.2270×10^4
eccentricity e	4.5×10^{-3}	1.4×10^{-2}	4.0×10^{-2}
inclination i (deg)	1.10×10^2	5.265×10^1	7.0×10^1

2 The Earth-LARES system

The Earth artificial satellite LARES/WEBER-SAT [6, 7] was proposed in order to measure the post-Newtonian general relativistic Lense-Thirring effect on the orbit of a test particle to a high accuracy level ($\sim 1\%$) in the gravitational field of the Earth by suitably combining its data with those of the existing geodetic SLR (Satellite Laser Ranging) satellites LAGEOS and LAGEOS II. Their orbital parameters are in Table 1.

2.1 The observational sensitivity

From $\Delta r = ea\Delta\omega$ [8], it can be obtained that the accuracy in measuring the perigee precession over a given observational time span can be expressed as $\delta\omega = \delta r/ea$. By assuming a root-mean-square (rms) error of 1 mm in reconstructing the LARES orbit over, say, one year for a given set of dynamical force models one gets $\delta\omega = 4 \times 10^{-1}$ mas.

With³ $a = 3.6 \times 10^4$ km and $e = 2.8 \times 10^{-1}$, as also suggested in [5], and by assuming a rms error of 1 cm, the accuracy in the perigee would amount to 2×10^{-1} mas. Note that the adopted values for the obtainable accuracies in r are optimistic; for example, the mm accuracy has not yet been fully achieved for the existing LAGEOS satellites.

2.2 The systematic errors of gravitational origin

The perigee of an Earth artificial satellite is affected by various kinds of long-period (i.e. averaged over one orbital revolution) orbital perturbations induced by the Newtonian Earth's geopotential [10]. The most insidious ones are the secular rates induced by the even ($\ell = 2, 4, 6, \dots$) zonal ($m = 0$) harmonic coefficients J_ℓ of the geopotential which account for the departure of the Earth from an exact spherical shape. Their explicit expressions up to

³It may be interesting to note that, with such an orbital configuration and $i = 6.34 \times 10^1$ deg the gravitomagnetic Lense-Thirring precession [?] would amount to -1.8 mas yr⁻¹ only.

degree $\ell = 20$ can be found, e.g., in [11]. The largest precession is induced by the Earth's quadrupole mass moment J_2 . For a moderate eccentricity its analytical expression is

$$\left. \frac{d\omega}{dt} \right|_{J_2} = \frac{3}{2} n \left(\frac{R}{a} \right)^2 \frac{J_2}{(1 - e^2)^2} \left(2 - \frac{5}{2} \sin^2 i \right), \quad (2)$$

where $n = \sqrt{GM/a^3}$ is the unperturbed Keplerian mean motion and R is the Earth's equatorial mean radius.

In [5] it is proposed to launch LARES in the so called frozen-perigee orbit characterized by the critical value of the inclination, $i = 6.34 \times 10^1$ deg, for which the J_2 -precession of eq.(2) vanishes. Moreover, it seems that Ciufolini suggests to use the LARES data together with those from LAGEOS and LAGEOS II in order to measure the Lense-Thirring as well.

We will now show that such proposals are unfeasible.

2.2.1 The impact of the even zonal harmonics of the geopotential

Indeed, apart from the fact that the unavoidable orbital injection errors would prevent to exactly insert LARES in orbit with the required inclination, it turns out that the impact of the other uncanceled precessions induced by the even zonal harmonics of higher degree, along with their secular variations, would totally swamp the LS effect. In Table 2 we use the calibrated sigmas of the even zonal harmonics of the recently released combined CHAMP+GRACE+terrestrial gravity EIGEN-CG01C Earth gravity model [12] in order to calculate the mismodelled residual classical precessions on the perigee of LARES by assuming $i = 6.34 \times 10^1$ deg. It can be easily seen that the mismodelled precessions induced by the first seven even zonal harmonics are larger than the LS precession. Due to the extreme smallness of such an effect, it is really unlikely that the forthcoming Earth gravity models from CHAMP and GRACE will substantially change the situation. This rules out the possibility of using only the perigee of LARES.

2.2.2 The linear combination approach

In regard to the possibility of suitably combining the Keplerian orbital elements of the existing LAGEOS satellites and of the proposed LARES [14, 15, 16] in order to reduce the impact of the systematic errors of the geopotential on the proposed measurement, it is unfeasible as well.

Indeed, the perigee is also affected, among other things, by the post-Newtonian general relativistic Einstein precession [17], whose nominal value

Table 2: Mismodelled classical secular precessions, in mas yr⁻¹, of the perigee of an Earth satellite with semimajor axis $a = 1.2270 \times 10^4$ km, inclination $i = 6.34 \times 10^1$ deg, eccentricity $e = 4.0 \times 10^{-2}$ according to the combined CHAMP+GRACE+terrestrial gravity EIGEN-CG01C Earth gravity model up to degree $\ell = 20$ (1-sigma). For the precessions induced by the secular variations of the even zonal harmonics, referred to an observational time span of one year, the values for $\dot{J}_\ell, \ell = 2, 4, 6$ of [13] have been used. Such precessions grow linearly in time. In the last three rows a - has been inserted because the corresponding mismodelled precessions amount to 10^{-4} mas yr⁻¹. The LS secular precession amounts to $\mp 4 \times 10^{-3}$ mas yr⁻¹ for an object orbiting a central mass along a circular path.

Even zonal harmonic	Mismodelled precessions (mas yr ⁻¹)
J_2	1.26×10^{-1}
$\dot{J}_2$	4×10^{-3} mas yr ⁻²
J_4	4.390
$\dot{J}_4$	1.406 mas yr ⁻²
J_6	1.355
$\dot{J}_6$	6.63×10^{-1} mas yr ⁻²
J_8	2.08×10^{-1}
J_{10}	2.5×10^{-2}
J_{12}	2.5×10^{-2}
J_{14}	7×10^{-3}
J_{16}	-
J_{18}	-
J_{20}	-

for LARES is $3.280136 \times 10^3 \text{ mas yr}^{-1}$, and by the Lense-Thirring effect, which, for $i = 6.34 \times 10^1 \text{ deg}$, nominally amounts to $-4.1466 \times 10^1 \text{ mas yr}^{-1}$. This means that if we want to measure the LS precession independently of such quite larger Newtonian and post-Newtonian effects we would need ten Keplerian orbital elements in order to write down a linear system of ten equations in ten unknowns (the first seven even zonal harmonics, the LS effect and the two relativistic precessions) and solve it for the LS precession. Instead, we would have at our disposal, in principle, only four reliable Keplerian orbital elements⁴: the longitudes of the ascending node Ω of LAGEOS, LAGEOS II and LARES-which are affected by the Earth's geopotential and by the Lense-Thirring effect but not by the LS force-and the perigee of LARES. They would only allow to cancel out the general relativistic effects and just one even zonal harmonic.

The data from an hypothetical satellite with $a = 3.6 \times 10^4 \text{ km}$, as proposed in [5], could not be used for the following reasons. If, on the one hand, the classical geopotential precessions would be smaller than the LS effect, apart from those induced by J_4 and J_6 according to EIGEN-CG01C, on the other hand, the gravitomagnetic Lense-Thirring and the gravitoelectric Einstein post-Newtonian precessions on the perigee would amount to $-1.8512 \text{ mas yr}^{-1}$ (for $i = 6.34 \times 10^1 \text{ deg}$) and $2.409956 \times 10^2 \text{ mas yr}^{-1}$, respectively. This means that, even for such a higher altitude, one could not analyze only the perigee of LARES whose Keplerian orbital elements should be, instead, combined with those of the existing LAGEOS and LAGEOS II: with the nodes of the LARES, LAGEOS and LAGEOS II and the perigee of LARES it would be possible, in principle, to disentangle the LS effect from the post-Newtonian precessions and one classical even zonal harmonic. For such a high altitude the period of the node of LARES would amount to $\sim 10^4$ days, i.e. tens of years. The tesseral K_1 tidal perturbation, which is one of the most powerful harmonic time-dependent perturbations which are not cancelled out by the linear combination approach, has just the period of the node. Then, it would act as a superimposed linear bias over an observational time span of a few years. Moreover, as shown in [18, 19], when high-altitude satellites are included in linear combinations involving also lower satellites as the existing LAGEOS and LAGEOS II it turns out that the orbital ele-

⁴Note that the other routinely and accurately laser-tracked SLR satellites which could, in principle, be considered are Ajsai, Starlette and Stella: the useful orbital elements are their nodes and the perigee of Starlette. However, since they orbit at much lower altitudes than the LAGEOS satellites they would practically be useless. Indeed, they are sensitive to much more even zonal harmonics of the geopotential so that they would greatly increase the systematic error induced by them.

ments of the higher SLR targets enter the combination with huge coefficients which amplify all the uncanceled orbital perturbations. This would also be the case for the K_1 tide affecting LARES.

2.2.3 The impact of the odd zonal harmonics

The perigee of an Earth artificial satellite is also affected by long-period harmonic perturbations induced by the odd ($\ell = 3, 5, 7, \dots$) zonal ($m = 0$) harmonics of the geopotential. The largest perturbation is induced by J_3 : it has a sinusoidal signature with the period of the perigee. Its analytic expression is [20]

$$\begin{aligned} \left. \frac{d\omega}{dt} \right|_{J_3} = & -\frac{3}{2}n \left(\frac{R}{a} \right)^3 \frac{J_3 \sin \omega}{e \sin i (1 - e^2)^3} \left[\left(\frac{5}{4} \sin^2 i - 1 \right) \sin^2 i + \right. \\ & \left. + e^2 \left(1 - \frac{35}{4} \sin^2 i \cos^2 i \right) \right]. \end{aligned} \quad (3)$$

For $i = 6.34 \times 10^1$ deg the period of the perigee of LARES, given by eq.(2), is of the order of 10^5 days; moreover, the second term of the right-hand-side of eq.(3) does not vanish. This means that the perigee of LARES in the critical inclination would be affected by an additional semisecular bias due to J_3 which, over an observational time span of some years, would resemble a superimposed linear trend. According to EIGEN-CG01C, its mismodelled effect would be $\leq 5 \times 10^{-1}$ mas yr $^{-1}$.

This additional bias should be accounted for both in the perigee-only scenario and in the linear combinations scenario. Note also that for $a = 3.6 \times 10^4$ km and $i = 6.34 \times 10^1$ deg the period of the perigee would amount to $\sim 10^7$ days, i.e. $\sim 10^4$ years.

3 The Earth-Moon system

In [1] the possibility of analyzing the perigee of the Moon ($a = 3.84 \times 10^5$ km, $e = 0.05490$) is considered in view of the possible future improvements in the range residuals accuracy from the cm to the mm level in the Lunar Laser Ranging (LLR) technique [21]. By assuming $\delta r \sim 1$ mm, the uncertainty in determining the Moon perigee would amount to $\delta\omega = 9 \times 10^{-3}$ mas which is the magnitude of the LS shift over almost 2 years. The currently available accuracy in LLR range residuals is ~ 2 cm which maps into a 1.8×10^{-1} mas accuracy on the perigee.

Table 3: Orbital parameters of Mercury, Venus and Mars (<http://nssdc.gsfc.nasa.gov/planetary/factsheet/>). For the Astronomical Unit (A.U.) we use the value 1 A.U.=149597870691 m. The angle ϵ refers to the inclination to the ecliptic: the inclination i to the reference plane is almost $\epsilon/2$.

Planet	a (A.U.)	ϵ (deg)	e
Mercury	3.8709893×10^{-1}	7.00487	2.0563069×10^{-1}
Venus	7.2333199×10^{-1}	3.39471	6.77323×10^{-3}
Mars	1.52366231	1.85061	9.341233×10^{-2}

4 The Solar System scenario

If we look at the planetary motions in the gravitational field of the Sun the situation is more favorable. The LS perihelion secular rates for all the planets, by neglecting the eccentricity of their orbits, amounts to⁵ $\mp 4 \times 10^{-4}$ " cy⁻¹. It is undoubtedly a small value if compared, e.g, to the well known Einstein perihelion advance which, for Mercury, amounts to 4.29811×10^1 " cy⁻¹.

4.1 The observational sensitivity

Contrary to the Earth-LARES scenario, the braneworld effect lies at the edge of the present-day available orbit determination accuracy.

The best tracked planet is Mars thanks to the various spacecraft missions to it: Viking (1976-1982), Pathfinder landers (1997), Mars Global Surveyor (MGS) (1998-2003) and Odyssey (2002-2003). According to [22], the rms residuals of ranging for Viking are 8 m, for Pathfinder are 4.4 m, for MGS and Odyssey are 1.4 m. From the results of Table 4 of [22] for the nonsingular elements h and k , based on the processing of about 300000 position observations (1911-2003) of different types used for the EPM2004 ephemerides, it turns out that the formal standard deviation in the Martian perihelion advance is 1.5×10^{-2} mas; the braneworld shift over a 92-years time span amounts to 3.68×10^{-1} mas.

The present-day situation for Mercury and, especially, Venus is less favorable mainly due to the systematic errors induced by the mismodelling

⁵Departures from such value due to the eccentricities are very likely too small to be detectable, as can be inferred from the subsequent discussion.

in the planetary topography and the small value of the eccentricity of the venusian orbit. Indeed, from [22] the rms residuals of ranging from all radiometric data (1911-2003) amount to 1.4 km and 0.7 km for Mercury and Venus, respectively. The formal standard deviations in the perihelia advance are 7.6×10^{-1} mas and 8.772 mas, respectively. However, the future Messenger (launched in August 2004 by NASA) BepiColombo and Venus Express (ESA) missions should improve, among other things, also the accuracy of the ephemerides of Mercury and Venus.

4.2 The systematic errors

However, as in the case of the Earth-LARES scenario, the systematic errors of gravitational origin represent a very limiting factor. In Table 4 the nominal values of the major Newtonian and post-Newtonian general relativistic competing precessions are listed. It turns out that the accuracy with which the various competing effects⁶ are presently known would make difficult to use only one perihelion, apart from, perhaps, Mars. The major limiting effects are the classical N-body precession, the post-Newtonian general relativistic gravitoelectric perihelion advance and the classical secular precession due to the Sun's even zonal harmonic J_2 .

A very limiting systematic bias might be, in principle, that induced by the huge Newtonian N-body secular precessions whose nominal values are up to seven orders of magnitude larger than the effect of interest. In regard to their mismodelling, the major source of uncertainty is represented by the GM of the perturbing bodies among which Jupiter plays a dominant role, especially for Mars. According to [25], the Jovian GM is known with a relative accuracy of 10^{-8} ; this would imply for the red planet a mismodelled precession induced by Jupiter of the order of 10^{-5} " cy^{-1} . The GM of Saturn is known with a 1×10^{-6} relative accuracy [26]. However, the ratio of the precession induced on the Mars perihelion by Saturn to that induced on the Mars perihelion by Jupiter is proportional to $(M_{\text{Sat}}/M_{\text{Jup}})(a_{\text{Jup}}/a_{\text{Sat}})^3 \sim 5 \times 10^{-2}$. This would assure that also the effect of Saturn is of the order of 10^{-5} " cy^{-1} . The situation with the precessions induced by Uranus and Neptune on the martian perihelion is even more favorable. Indeed, for Uranus $(M_{\text{Ura}}/M_{\text{Jup}})(a_{\text{Jup}}/a_{\text{Ura}})^3 \sim 1 \times 10^{-3}$ and the relative uncertainty in the uranian GM is 2×10^{-6} [27]. For Neptune $(M_{\text{Nep}}/M_{\text{Jup}})(a_{\text{Jup}}/a_{\text{Nep}})^3 \sim 3 \times 10^{-4}$ and $\delta(GM)/GM = 2 \times 10^{-6}$ [28]. In regard to the precessions induced by the most important asteroids, their

⁶Note that the nominal values of the Lense-Thirring precessions are one order of magnitude smaller than the LS effect, so that the gravitomagnetic force does not pose problems.

Table 4: Nominal values, in $" \text{ cy}^{-1}$, of the Newtonian and post-Newtonian general relativistic secular precessions affecting the perihelia and the nodes of the inner planets of the Solar System. For a given planet, the precession labelled with N-body is due to the classical interactions with the other bodies of the Solar System, that labelled with GE is due to the post-Newtonian gravitoelectric Schwarzschild component of the solar gravitational field, that labelled with J_2 is due to the classical effect of the Sun's quadrupole mass moment J_2 and that labelled with LT is due to the post-Newtonian gravitomagnetic Lense-Thirring component of the solar gravitational field. The LS perihelion secular rates for all the planets, by neglecting the eccentricities of the orbits, amount to $3.95 \times 10^{-4} "$ cy^{-1} . For the numerical calculations of the planetary N-body precessions see <http://ssd.jpl.nasa.gov/elemp/planets.html#rates>. For the Sun's proper angular momentum J , which is the source of the gravitomagnetic field, the value $1.9 \times 10^{41} \text{ kg m}^2 \text{ s}^{-1}$ [23] has been adopted. For the Sun-planetary data see [24].

Secular rate	Mercury	Venus	Mars
$\dot{\omega}_{\text{N-body}}$	1.019036×10^3	8.87652×10^2	2.580836×10^3
$\dot{\Omega}_{\text{N-body}}$	-4.4630×10^2	-9.9689×10^2	-1.02019×10^3
$\dot{\omega}_{\text{GE}}$	4.2981×10^1	8.624	1.351
$\dot{\Omega}_{\text{GE}}$	0	0	0
$\dot{\omega}_{J_2}$	5.0609×10^{-2}	5.223×10^{-3}	3.92×10^{-4}
$\dot{\Omega}_{J_2}$	-2.5375×10^{-2}	-2.613×10^{-3}	-1.96×10^{-4}
$\dot{\omega}_{\text{LT}}$	-3.018×10^{-3}	-4.31×10^{-4}	-4.5×10^{-5}
$\dot{\Omega}_{\text{LT}}$	1.008×10^{-3}	1.44×10^{-4}	1.5×10^{-5}

Table 5: First row: Formal standard deviations, in $" \text{ cy}^{-1}$, in the measurement of the post-Newtonian secular rates of the planetary perihelia from the processing of the ranging observations (E.V. Pitjeva, private communication, 2004). They have been obtained by including also the perturbing effects of the asteroids which are particularly important for Mars (see Appendix 6). The uncertainty in the perihelion advances is one-to-three orders of magnitude larger than the LS effect for Mercury and Venus, respectively, but it is slightly smaller than the LS precession for Mars. Realistic errors may be an order of magnitude larger. Second row: Formal standard deviations, in $" \text{ cy}^{-1}$, in the numerical propagation of the planetary nodal rates averaged over 200 years from the DE405 ephemerides (E.M. Standish, private communication, 2004).

Secular rate	Mercury	Venus	Mars
$\dot{\omega}$	8.6×10^{-3}	1.037×10^{-1}	1×10^{-4}
$\dot{\Omega}$	1.82×10^{-4}	6×10^{-6}	1×10^{-6}

masses are not known with great accuracy [29]; according to recent estimations, their mismodelled effect would amount to $\sim 10^{-4} " \text{ cy}^{-1}$ for the secular terms (see Appendix 6).

In regard to the post-Newtonian relativistic perihelion advance, it is currently known at a 10^{-4} level of relative accuracy from the PPN parameters β and γ [22]. It turns out that the accuracy with which the post-Newtonian general relativistic gravitoelectric perihelion advances can be determined is one-to-three orders of magnitude larger than the LS effect for Mercury and Venus, respectively, but it is slightly smaller than the LS precession for Mars. The figures of Table 5 for the perihelia have been obtained from the processing of about 300000 position observations (1911-2003) of different types: also the perturbing effects of the asteroids [22], which are particularly important for Mars (see Appendix 6), have been accounted for. However, it must be noted that the realistic errors in such figures may be one order of magnitude larger, so that the mismodelling in the relativistic effect would tend to mask the LS rate.

The mismodelled precessions induced by J_2 , by assuming a relative uncertainty of 10% in it [22], would be larger than the LS effect for Mercury and Venus; for Mars such competing classical feature would, instead, be smaller than the LS effect by one order of magnitude.

These evaluations would suggest that in a more or less near future it might, perhaps, be possible to analyze the perihelion of Mars only. However, note that a sort of a priori ‘imprint’ of the LS signature in the recovery of all the other effects affecting the perihelion of Mars would be present: it might make unreliable the proposed measurement by driving the outcome of the test just towards the desired result. Then, a better approach would consist in disentangling the LS effect from all the other competing effects. This can be done by suitably combining the perihelion of Mars with other Keplerian orbital elements affected by such Newtonian and post-Newtonian precessions. A similar approach has recently been adopted, in a Solar System context, in [30, 31]. By writing down the rates of the perihelia and the nodes of Mars and Mercury in terms of the LS, general relativistic, J_2 and N-Body precessions it is possible to consider them as a linear nonhomogeneous algebraic system of four equations in four unknowns and to solve it with respect to the LS effect. The so obtained linear combination of the residuals of the rates of the perihelia and the nodes of Mercury and Mars is, by construction, independent of the effects induced by the Newtonian N-body and solar precessions and by the general relativistic gravitoelectric force. The result is

$$\delta\dot{\omega}^{\text{Mars}} + c_1\delta\dot{\Omega}^{\text{Mars}} + c_2\delta\dot{\omega}^{\text{Mercury}} + c_3\delta\dot{\Omega}^{\text{Mercury}} = X_{\text{Dv}}, \quad (4)$$

where

$$c_1 = 2.5275, \quad c_2 = -3.14 \times 10^{-2}, \quad c_3 = -6.67 \times 10^{-2}, \quad (5)$$

and

$$X_{\text{Dv}} = \mp 3.87 \times 10^{-4} \text{ '' cy}^{-1}. \quad (6)$$

It is important to note that the impact of Mercury is relatively small, as can be inferred from eq.(5). Note that eq.(4) does not cancel out the Lense-Thirring effect: however, according to Table 4, its nominal impact amounts to $2 \times 10^{-5} \text{ '' cy}^{-1}$. According to the results for the nonsingular elements h , k , p , q of Table 4 of [22], The formal 1-sigma error affecting eq.(4) can be evaluated as 41%.

5 Conclusions

In this paper we have investigated the possibility of testing the multidimensional brane-world scenario proposed by Dvali, Gabadadze and Porrati by analyzing in the weak-field regime of the Solar System the pericentre advance derived by Lue and Starkman.

It has been shown that the systematic errors induced by the multipolar expansion of the classical part of the terrestrial gravitational potential rule out the possibility, suggested by Ciufolini, of using the perigee of the proposed LARES/WEBER-SAT satellite in order to test the Dvali, Gabadadze and Porrati gravity by measuring the Lue-Starkman pericentre precession. Indeed, according to the EIGEN-CG01C Earth gravity model from the dedicated CHAMP and GRACE missions, the errors induced by the mismodelled even zonal harmonics of the multipolar expansion of the Earth's gravitational potential would be up to three orders of magnitude larger than the LS precession of interest which amounts to $\mp 4 \times 10^{-3}$ mas yr $^{-1}$ only. Moreover, also the observational accuracy, evaluated with optimistic assumptions, would not be sufficient to measure such an effect with the LARES perigee.

On the contrary, from the ranging data to Mars and Mercury it might be possible, in a near future, to try to test the LS precession, which amounts to $\mp 4 \times 10^{-1}$ mas cy $^{-1}$, with a suitable linear combination of the residuals of the perihelia and the nodes of Mars and Mercury. Indeed, the perihelion advance of Mars, which is the currently best tracked planet of the Solar System, has recently been measured with a formal standard deviation of 1.5×10^{-2} mas over a time span of 92 years. The main systematic errors of gravitational origin, which severely affects a single perihelion rate, are instead cancelled out by the proposed linear combination. The obtainable accuracy could roughly be evaluated to be of the order of 41% at 1-sigma.

6 Appendix: The impact of the asteroids on the Mars orbital motion

The bodies of the asteroid's belt induce both secular and long-period perturbations on the orbital elements of Mars whose amplitudes are proportional, among other things, to their masses. Since they are relatively poor determined with respect to those of the major planets of the Solar System the mismodelled part of the orbital perturbations induced by them might become relevant also because their nominal size is larger than the LS effect. It is important to note that the periods of some of the most powerful perturbations amount to tens and even hundreds of years: it would be a major limiting factor for analysis covering relatively short time spans of available data.

In the framework of the linear Lagrange-Laplace perturbation theory the

equation for the rate of the longitude of the perihelion is

$$\frac{d\varpi}{dt} = \frac{\sqrt{1-e^2}}{na^2e} \frac{\partial \mathcal{R}}{\partial e} + \frac{\tan(i/2)}{na^2\sqrt{1-e^2}} \frac{\partial \mathcal{R}}{\partial i}, \quad (7)$$

where $\mathcal{R}$ is the disturbing function which accounts for the effect of a third body, whose parameters are denoted with $'$, on a given planet, whose parameters are denoted without $'$. It is

$$\mathcal{R} = GM' \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} - \frac{\mathbf{r} \cdot \mathbf{r}'}{r'^3} \right). \quad (8)$$

Eq.(8) can be expanded as a multiple Fourier series with respect to the six orbital parameters of the perturbed and perturbing bodies $\lambda, \varpi, \Omega, \lambda', \varpi', \Omega'$, where λ is the mean longitude

$$\mathcal{R} = \sum_{jj' ll' mm'} F_{jj' ll' mm'} \cos(j\lambda + j'\lambda' + l\varpi + l'\varpi' + m\Omega + m'\Omega'). \quad (9)$$

In turn, the coefficients $F_{jj' ll' mm'}$, which depend on a, a', e, e', i, i' , can be expanded as a Taylor series with respect to $e, e', \sin i, \sin i'$ due to the smallness of the eccentricities and the inclinations of the Solar System bodies

$$F_{jj' ll' mm'} = \sum_{r_1=0}^{\infty} \sum_{r_2=0}^{\infty} \sum_{r_3=0}^{\infty} \sum_{r_4=0}^{\infty} C_{r_1 r_2 r_3 r_4} e^{|l|+2r_1} e'^{|l'|+2r_2} \sin^{|m|+2r_3} i \sin^{|m'|+2r_4} i'. \quad (10)$$

In it the coefficients $C_{r_1 r_2 r_3 r_4}$ depend on a and a' . In eq.(9) the so called short-period term are those which contain the mean longitudes λ and λ' , while the secular terms are those which do not contain them, i.e. for which $j = j' = 0$. In regard to the latter ones, it turns out that $\langle \mathcal{R} \rangle$, averaged over the mean longitudes, only contain the terms $1, e^2, e'^2, \sin^2 i, \sin^2 i', ee' \cos(\varpi - \varpi'), \sin i \sin i' \cos(\Omega - \Omega')$.

Let us now specialize these general concepts to the Mars-asteroids system.

- According to [32], the nominal amplitude of the secular perturbation on the longitude of the perihelion of Mars induced by the major asteroids amount to $3 \times 10^{-2}'' \text{ cy}^{-1}$. The three most massive bodies (1) Ceres, (2) Pallas and (4) Vesta are responsible of roughly 80% of such perturbation, i.e. $\sim 2.4 \times 10^{-2}'' \text{ cy}^{-1}$. The latest reported measurements of the masses of the asteroids [22] are precise up to $\sim 0.07-0.3\%$,

so that the mismodelled part of the Mars perihelion secular precession due to the three major asteroids amounts to $\sim 1 \times 10^{-4}'' \text{ cy}^{-1}$. The masses of the smaller asteroids like (3) Juno, (7) Iris and (324) Bamberga are known to a $\sim 1\%$ level, so that the mismodelled part of their perturbation may be evaluated as $\sim 2 \times 10^{-4}'' \text{ cy}^{-1}$. The impact of the asteroid ring, i.e. the ensemble of the minor asteroids which can be modelled as due to a solid ring in the ecliptic plane [33], can be worked out, e.g., with the Lagrangian approach and the disturbing function of the Appendix of [33]. By using the values of [22] for the ring's radius and mass it turns out that the secular perturbation on $\dot{\omega}_{\text{Mars}}$ amounts to $2.8 \cdot 10^{-3}'' \text{ cy}^{-1}$, with an uncertainty of $3 \cdot 10^{-4}'' \text{ cy}^{-1}$.

- In regard to the short-period perturbations, according to [32], the terms of eq.(9) containing also the mean longitudes are, in principle, not negligible because of the occurrence of many orbital resonances which induce harmonic perturbations with periods tens or even hundreds of years long. The focus in [32] is on the perturbations of the mean longitudes: the nominal values of their amplitudes are of the order of $10^{-3} - 10^{-4}'' \text{ cy}^{-1}$ (see Table IV of [32]). The current accuracy in our knowledge of the asteroids masses makes the mismodelled part of such perturbations smaller than the investigated braneworld effect: moreover, it should also be pointed out that the perturbations in the mean longitudes, which depend on the square of their periods, are larger than those in the other orbital elements, which, instead, depend linearly on their periods. These considerations would suggest that the resonant harmonic perturbations on the Martian perihelion due to the asteroids should not compromise the recovery of the LS effect. However, a careful and more detailed analysis of such an important topic is needed.

Concerning the error in the observational determination of the Martian perihelion rate, according [32] in the DE 118 and DE 200 JPL ephemerides it was $6 \times 10^{-3}'' \text{ cy}^{-1}$; with the recent EPM2004 IAA (RAS) ephemerides it amounts to $1 \times 10^{-4}'' \text{ cy}^{-1}$ (E.V. Pitjeva, personal communication 2004).

Acknowledgements

I am grateful to E. V. Pitjeva and E. M. Standish for their help and clarifications. Thanks also to G. Starkman for his useful remarks, to T.M. Eubanks

for the important material sent to me and for relevant discussions on the impact of the asteroids on the Martian orbit and to J. Williams for his clarifications on the possibilities of LLR.

References

- [1] Dvali, G., Gruzinov, A., and Zaldarriaga, M. (2003). The Accelerated Universe and the Moon, *Phys. Rev. D*, **68** 024012.
- [2] Dvali, G., Gabadadze, G., and Porrati, M. (2000). 4D Gravity on a Brane in 5D Minkowski Space, *Phys. Lett. B*, **485** 208–214.
- [3] Lue, A., and Starkman, G. (2003). Gravitational Leakage into Extra Dimensions Probing Dark Energy Using Local Gravity, *Phys. Rev. D*, **67** 064002.
- [4] Dvali, G. (2004). Infrared Modification of Gravity, Preprint hep-th/0402130.
- [5] Ciufolini, I. (2004). LARES/WEBER-SAT, frame-dragging and fundamental physics, Preprint gr-qc/0412001.
- [6] Ciufolini, I. (1986). Measurement of the Lense-Thirring Drag on High-Altitude, Laser Ranged Artificial Satellite, *Phys. Rev. Lett.*, **56** 278-281.
- [7] Iorio, L., Lucchesi, D., and Ciufolini, I. (2002). The LARES Mission Revisited: An Alternative Scenario, *Class. Quantum Grav.*, **19** 4311-4325.
- [8] Nordtvedt, K. (2000). Improving gravity theory tests with solar system “grand fits”, *Phys. Rev. D*, **61**, 122001.
- [9] Lense, J., and Thirring, H. (1918). Über den Einfluss der Eigenrotation der Zentralkörper auf die Bewegung der Planeten und Monde nach der Einsteinschen Gravitationstheorie, *Phys. Z.*, **19**, 156-163. Translated and discussed by Mashhoon, B., Hehl, F.W., and Theiss, D.S. (1984). On the Gravitational Effects of Rotating Masses: The Thirring-Lense Papers, *Gen. Rel. Grav.*, **16**, 711-750.
- [10] Kaula, W. M. (1966). *Theory of Satellite Geodesy* (Waltham: Blaisdell Publishing Company).

- [11] Iorio, L. (2003). The Impact of the Static Part of the Earth's Gravity Field on Some Tests of General Relativity with Satellite Laser Ranging, *Celest. Mech. & Dyn. Astron.*, **86** 277-294.
- [12] Reigber, Ch., Schwintzer, P., Stubenvoll, R., Schmidt, R., Flechtner, F., Meyer, U., Knig, R., Neumayer, H., Frste, Ch., Barthelmes, F., Zhu, S.Y., Balmino, G., Biancale, R., Lemoine, J.-M., Meixner, H., and Raimondo, J.C. (2005). A High Resolution Global Gravity Field Model Combining CHAMP and GRACE Satellite Mission and Surface Gravity Data: EIGEN-CG01C, *J. of Geodesy*, at press.
- [13] Cox, C., Au, A., Boy, J.-P., and Chao, B. (2003). Time-Variable Gravity: Using Satellite Laser Ranging as a Tool for Observing Long-Term Changes in the Earth System. *Proc. 13th Int. Laser Ranging Workshop NASA CP 2003-212248*, R. Noomen, S. Klosko, C. Noll, and M. Pearlman (Eds.), (NASA Goddard) (Preprint http://cddisa.gsfc.nasa.gov/lw13/lw_proceedings.html#science).
- [14] Ciufolini, I. (1996). On a new method to measure the gravitomagnetic field using two orbiting satellites, *Il Nuovo Cimento A*, **109**, 1709-1720.
- [15] Iorio, L. (2004). The impact of the new CHAMP and GRACE Earth gravity models on the measurement of the general relativistic Lense-Thirring effect with the LAGEOS and LAGEOS II satellites, *Earth Observation with CHAMP - Results from Three Years in Orbit* ed Ch. Reigber, H. Lüher, P. Schwintzer, and J. Wickert (Springer, Berlin) pp 187–192.
- [16] Iorio, L., and Doornbos, E. (2005). On the possibility of using Jason-1 in determining the Lense-Thirring effect, *Gen. Rel. Grav.*, in press. Preprint gr-qc/0404062.
- [17] Einstein, A. (1915). Erklärung der perihelbewegung des Merkur aus der allgemeinen Relativitätstheorie, *Sitzber. Kön. Preuss. Akad. Berlin*, **47**, 831-839.
- [18] Iorio, L. (2002). Is it possible to improve the present LAGEOS-LAGEOS II Lense-Thirring experiment?, *Class. Quantum Grav.* **19**, 5473-5480.

- [19] Vespe, F., and Rutigliano, P. (2004). The improvement of the Earth gravity field estimation and its benefits in the atmosphere and fundamental physics, paper presented at *35th COSPAR Scientific Assembly, Paris, France, 18 - 25 July 2004*, COSPAR04-A-03614.
- [20] Kiedron, K., and Cook, R. (1991). Frozen Orbits in the J2+J3 Problem, paper presented at *AAS/AIAA Astrodynamics Specialist Conference, Durango, Colorado, 19-22 August 1991*, AAS 91-426.
- [21] Turyshev, S.G., Williams, J.G., Nordtvedt, K., Shao, M., and Murphy, T.W. (2004). 35 Years of Testing Relativistic Gravity: Where Do We Go from Here? *Astrophysics, Clocks and Fundamental Constants* ed Karshenboim, S.G., and Peik, E. (Springer, Berlin) pp 311-330.
- [22] Pitjeva, E.V. (2004b). Precise determination of the motion of planets and some astronomical constants from modern observations, to appear in *Proceedings Transit of Venus: New Views of the Solar System and Galaxy IAU Colloquium No. 196, 2004* ed D.W. Kurtz, and G.E. Bromage, (Cambridge University Press, Cambridge).
- [23] Pijpers, F. P. (2003). Astroseismic determination of stellar angular momentum, *Astron. and Astrophys.*, **402**, 683-692.
- [24] Seidelmann, K.P. (ed.) (1992). *Explanatory Supplement to the Astronomical Almanac*, (University Science Books, Mill Valley, California), p 316 (Table 5.8.1).
- [25] Jacobson, R.A. (2003). JUP230 orbit solution.
- [26] Jacobson, R.A. (2004). The orbits of the major saturnian satellites and the gravity field of Saturn from spacecraft and Earth-based observations, *Astron. J.*, **501**, 492-501.
- [27] Jacobson, R. A., Campbell, J. K., Taylor, A. H., and Synnott, S. P. (1992). The masses of Uranus and its major satellites from Voyager tracking data and Earth-based Uranian satellite data, *Astron. J.*, **103**, 2068-2078.
- [28] Jacobson, R. A., Riedel, J. E., and Taylor, A. H. (1991). The orbits of Triton and Nereid from spacecraft and Earthbased observations, *Astron. and Astroph.*, **247**, 565-575.

- [29] Standish, E.M. (2002). *Proceedings of the IERS Workshop on the Implementation of the New IAU Resolutions, IERS Technical Note No. 29* ed N. Capitaine et al, (Verlag des Bundesamts für Kartographie und Geodäsie, Frankfurt am Main), p 61.
- [30] Iorio, L. (2005a). On the possibility of measuring the solar oblateness and some relativistic effects from planetary ranging, *Astron. and Astroph.*, **433**, 385-393.
- [31] Iorio, L. (2005b). Is it possible to measure the Lense-Thirring effect on the orbits of the planets in the gravitational field of the Sun? , *Astron. and Astroph.*, **431**, 385-389.
- [32] Williams, J.G. (1984). Determining Asteroid Masses from Perturbations on Mars, *Icarus*, **57**, 1-13.
- [33] Krasinsky, G.A., Pitjeva, E.V., Vasiljev, M.V., and Yagudina, E.I. (2002). Hidden Mass in the Asteroid Belt, *Icarus*, **158**, 98-105.